

COHOMOLOGY IN ONE-DIMENSIONAL SUBSTITUTION TILING SPACES

MARCY BARGE AND BEVERLY DIAMOND

ABSTRACT. Anderson and Putnam showed that the cohomology of a substitution tiling space may be computed by collaring tiles to obtain a substitution which “forces its border.” One can then represent the tiling space as an inverse limit of an inflation and substitution map on a cellular complex formed from the collared tiles; the cohomology of the tiling space is computed as the direct limit of the homomorphism induced by inflation and substitution on the cohomology of the complex. For one-dimensional substitution tiling spaces, we describe a modification of the Anderson-Putnam complex on collared tiles that allows for easier computation and provides a means of identifying certain special features of the tiling space with particular elements of the cohomology.

2000 Mathematics Subject Classification: *Primary:* 37B05; *Secondary:* 54H20, 55N05

1. INTRODUCTION

The investigation of the topology of substitution tiling spaces, in particular, the study of the cohomology of the space, is often aided by representing the tiling space as the inverse limit of an inflation and substitution map on the cellular Anderson-Putnam complex ([AP]). If the substitution *forces its border*, the complex can be formed using the given tiles as cells, and adding identifications indicating allowed transitions; the cohomology of the tiling space is computed as the direct limit of the homeomorphism induced by inflation and substitution on the cohomology of the complex. If the substitution does not force its border, then the inverse limit constructed in this way may be a nontrivial quotient of the tiling space rather than the tiling space itself; certain arc components may be glued together, and others pinched at points, in some cases changing the cohomology. In this case, the process of *collaring* provides a larger set of tiles and an induced substitution which does force its border. The inverse limit space of the complex formed from the larger set of tiles and associated map is homeomorphic to the tiling space of the original substitution.

In many situations, full collaring leads to a very large number of tiles, and the process of determining the complex can be tedious and time-consuming.

We indicate below, for one-dimensional substitution tilings (that is, the tiles are subintervals of $\mathbb{R}$), a process which achieves the same ultimate goal, that of representing the tiling space as an inverse limit on an associated, simple complex, from which the cohomology can be computed in a fairly straightforward way.

One can always *rewrite* a one-dimensional substitution to obtain a *proper* substitution (see [Dur] and [BD]) that then automatically forces its border. The topology of the tiling space is unchanged by rewriting ([BD]), and the resulting proper substitution typically has a smaller alphabet than that of the substitution produced by collaring, hence cohomology may be more efficiently computed by rewriting than by collaring. This process still has two drawbacks in our view. First, there is no analog of rewriting in higher dimensions, while the construction we describe below generalizes easily to higher dimensions. Second, the approach we present yields a specific, concrete, topological interpretation of a piece of the cohomology.

2. NOTATION AND TERMINOLOGY

We introduce some notation and terminology.

Let $\mathcal{A} = \{1, 2, \dots, \text{card}(\mathcal{A})\}$ be a finite alphabet; $\mathcal{A}^*$ will denote the collection of finite nonempty words with letters in $\mathcal{A}$. A *substitution* on $\mathcal{A}$ is a map $\varphi : \mathcal{A} \rightarrow \mathcal{A}^*$; φ extends naturally to $\varphi : \mathcal{A}^* \rightarrow \mathcal{A}^*$. The *transition matrix* $A_\varphi = A$ for φ is $A := (a_{ij})_{i \in \mathcal{A}, j \in \mathcal{A}}$ in which a_{ij} is the number of occurrences of i in the word $\varphi(j)$.

The substitution φ is *primitive* if $\varphi^n(i)$ contains j for all $i, j \in \mathcal{A}$ and sufficiently large n . Equivalently, φ is primitive if and only if the matrix A is aperiodic, in which case A has a simple eigenvalue λ_φ larger in modulus than its remaining eigenvalues called the Perron-Frobenius eigenvalue of A (and φ).

A word w is *allowed* for φ if and only if for each finite subword (i.e., factor) w' of w , there are $i \in \mathcal{A}$ and $n \in \mathbb{N}$ such that w' is a subword of $\varphi^n(i)$; the *language* of φ , $\mathcal{L}_\varphi = \mathcal{L}$, is the set of finite allowed words for φ . Let W_φ denote the set of allowed bi-infinite words for φ . We identify the 0^{th} coordinate in a bi-infinite word w by either an indexing, as in $w = \dots w_{-1}w_0w_1\dots$, or by use of a decimal point (or both). Let $\sigma : W_\varphi \rightarrow W_\varphi$ denote the *shift map*:

$$\sigma(\dots w_{-1}.w_0w_1\dots) := \dots w_{-1}w_0.w_1\dots$$

The substitution $\varphi : \mathcal{A} \rightarrow \mathcal{A}^*$ extends to $\varphi : W_\varphi \rightarrow W_\varphi$ where

$$\varphi(\dots w_{-1}w_0w_1\dots) := \dots \varphi(w_{-1}) \cdot \varphi(w_0)\varphi(w_1)\dots$$

The word w is *periodic* for φ under inflation and substitution, or *φ -periodic*, if for some $m \in \mathbb{N}$,

$$\varphi^m(w) = \dots \varphi^m(w_{-1}).\varphi^m(w_0)\varphi^m(w_1)\dots = \dots w_{-1}.w_0w_1\dots$$

Each primitive substitution φ has at least one allowed φ -periodic bi-infinite word which is necessarily uniformly recurrent under the shift. A substitution φ with precisely one periodic, hence fixed, bi-infinite word is called *proper*; φ is proper if and only if there are $b, e \in \mathcal{A}$ such that for all sufficiently large k and all $i \in \mathcal{A}$, $\varphi^k(i) = b \dots e$.

A primitive substitution φ is *aperiodic* if at least one (equivalently, each) φ -periodic bi-infinite word is not periodic under the natural shift map, in which case (W_φ, σ) is an infinite minimal dynamical system. If φ is aperiodic, then the map $\varphi : W_\varphi \rightarrow W_\varphi$ is one-to-one ([Mo]). If φ is periodic (that is, primitive and not aperiodic), then W_φ is finite.

Given a primitive substitution $\varphi : \mathcal{A} \rightarrow \mathcal{A}^*$ with $\text{card}(\mathcal{A}) = d \geq 2$, let $\omega_L := (\omega_1, \dots, \omega_d)$ be a positive left eigenvector for the Perron-Frobenius eigenvalue, λ , of A . The intervals $P_i = [0, \omega_i]$, $i = 1, \dots, d$, are called *prototiles* for φ (consider P_i to be distinct from P_j for $i \neq j$ even if $\omega_i = \omega_j$). A *tiling* T of $\mathbb{R}$ by the prototiles for φ is a collection $T = \{T_i\}_{i=-\infty}^\infty$ of tiles T_i for which $\bigcup_{i=-\infty}^\infty T_i = \mathbb{R}$, each T_i is a translate of some P_j (in which case we say T_i is of *type* j), and $T_i \cap T_{i+1}$ is a singleton for each i . Generally we assume that the indexing is such that $0 \in T_0 \setminus T_1$.

If $\varphi(i) = i_1 i_2 \dots i_{k(i)}$, then $\lambda \omega_i = \sum_{j=1}^{k(i)} \omega_{i_j}$. Thus $|\lambda P_i| = \sum_{j=1}^{k(i)} |P_{i_j}|$, and λP_i is tiled by $\{T_j\}_{j=1}^{k(i)}$, where $T_j = P_{i_j} + \sum_{k=1}^{j-1} \omega_{i_k}$. This process is called *inflation and substitution* and extends to a map Φ taking a tiling $T = \{T_i\}_{i=-\infty}^\infty$ of $\mathbb{R}$ by prototiles to a new tiling, $\Phi(T)$, of $\mathbb{R}$ by prototiles defined by inflating, substituting, and suitably translating each T_i . More precisely, for $w = w_1 \dots w_n \in \mathcal{A}^*$, define

$$\mathcal{P}_w + t = \{P_{w_1} + t, P_{w_2} + t + |P_{w_1}|, \dots, P_{w_n} + t + \sum_{i < n} |P_{w_i}|\}.$$

Then $\Phi(P_i + t) = \mathcal{P}_{\varphi(i)} + \lambda t$ and $\Phi(\{P_{k_i} + t_i\}_{i \in \mathbb{Z}}) = \bigcup_{i \in \mathbb{Z}} (\mathcal{P}_{\varphi(k_i)} + \lambda t_i)$.

There is a natural topology on the collection Σ_φ of all tilings of $\mathbb{R}$ by prototiles ($\{T_i\}_{i=-\infty}^\infty$ and $\{T'_i\}_{i=-\infty}^\infty$ are “close” if there is an ϵ near 0 so that $\{T_i\}_{i=-\infty}^\infty$ and $\{T'_i + \epsilon\}_{i=-\infty}^\infty$ are identical in a large neighborhood of 0 (see [AP] for details)). The space Σ_φ is compact and metrizable with this topology and $\Phi : \Sigma_\varphi \rightarrow \Sigma_\varphi$ is continuous. Given $T = \{T_i\}_{i=-\infty}^\infty \in \Sigma_\varphi$, let $\underline{w}(T) = \dots w_{-1} w_0 w_1 \dots$ denote the bi-infinite word with $w_i = j$ if and only if T_i is of type j . The *tiling space associated with* φ , $\mathcal{T}_\varphi$, is defined as

$$\mathcal{T}_\varphi = \{T : \underline{w}(T) \text{ is allowed for } \varphi\}.$$

There is a natural flow (translation) on Σ_φ defined by $(\{T_i\}_{i=-\infty}^\infty, t) \mapsto \{T_i - t\}_{i=-\infty}^\infty$. If φ is primitive and aperiodic, $\Phi : \mathcal{T}_\varphi \rightarrow \mathcal{T}_\varphi$ is a homeomorphism (this relies on the notion of *recognizability* or invertibility for such substitutions—see [Mo] and [So]). Each $T \in \mathcal{T}_\varphi$ is uniformly recurrent under the flow and has dense orbit (i.e., the flow is minimal on $\mathcal{T}_\varphi$). It follows that $\mathcal{T}_\varphi$ is a continuum.

Recall that a *composant* of a point x in a topological space X is the union of the proper compact connected subsets of X containing x . If φ is a primitive substitution, composants and arc components in $\mathcal{T}_\varphi$ are identical; in this case we use the terms interchangeably. For any substitution φ , the arc components of the tiling space $\mathcal{T}_\varphi$ coincide with the orbits of the natural flow (translation) on $\mathcal{T}_\varphi$.

Tilings $T, T' \in \mathcal{T}_\varphi$ are *forward asymptotic* if $\lim_{t \rightarrow \infty} \text{dist}(T - t, T' - t) = 0$. Equivalently, $T = \{T_i\}_{i=-\infty}^\infty, T' = \{T'_i\}_{i=-\infty}^\infty$ are *forward asymptotic* if there are $N, M \in \mathbb{Z}$ so that $T_{N+k} = T'_{M+k}$ for all $k \geq 0$. Composants are *forward asymptotic* if they contain forward asymptotic tilings. Backward asymptotic tilings and composants are defined similarly.

If $f : X \rightarrow X$ is a map of a compact connected metric space X , then the *inverse limit space* with single bonding map f is the space

$$\varprojlim f = \{(x_0, x_1, \dots) : f(x_i) = x_{i-1} \text{ for } i = 1, 2, \dots\}$$

with metric

$$d(\underline{x}, \underline{y}) = \sum_{i \geq 0} \frac{d(x_i, y_i)}{2^i};$$

$\hat{f} : \varprojlim f \rightarrow \varprojlim f$ will denote the natural (shift) homeomorphism

$$\hat{f}(x_0, x_1, \dots) = (f(x_0), x_0, x_1, \dots).$$

3. THE MODIFIED COMPLEX AND ITS COHOMOLOGY

Given a primitive, aperiodic substitution φ on d letters, the complex $K = K_\varphi$ will consist of a collection of edges representing the letters of $\mathcal{A}$, another collection of edges representing allowed transitions between letters, and certain identifications.

As above, $\omega_L = (\omega_1, \dots, \omega_d)$ is a left eigenvector for the Perron-Frobenius eigenvalue λ for φ . Define $0 < \epsilon = \min\{\frac{\omega_a}{2\lambda}\}_{a \in \mathcal{A}}$. For $a \in \mathcal{A}$, let

$$e_a := [\epsilon, \omega_a - \epsilon] \times \{a\},$$

and for $ab \in \mathcal{L}$, let

$$e_{ab} := [-\epsilon, \epsilon] \times \{ab\}.$$

Define

$$K := [(\cup_{a \in \mathcal{A}} e_a) \cup (\cup_{ab \in \mathcal{L}} e_{ab})] / \sim,$$

where

$$(\omega_a - \epsilon, a) \sim (-\epsilon, ab)$$

and

$$(\epsilon, b) \sim (\epsilon, ab)$$


for all $a, b \in \mathcal{A}$, and let

$$S := \cup_{ab \in \mathcal{L}} e_{ab} / \sim.$$

Loosely, the complex K is a wedge of d circles with the branch point ‘blown up’ to S ; K/S is homeomorphic to the wedge of d circles.

Example 1. *The Fibonacci substitution:*

If φ is the Fibonacci substitution ($\varphi(1) = 12, \varphi(2) = 1$), then all transitions except 22 are allowed, and K_φ is (topologically and not to scale) shown below.



(In this case, collaring leads to a topologically identical complex.) $\square$

Given a tiling $T = \{T_n\}_{n \in \mathbb{Z}} \in \mathcal{T}_\varphi$, let $t \in \mathbb{R}$ and $a, b, c \in \mathcal{A}$ be such that $0 \in T_0 = [0, \omega_a] - t$, $T_{-1} = [0, \omega_b] - t - \omega_b$, and $T_1 = [0, \omega_c] - t + \omega_a$. Define a map $p : \mathcal{T}_\varphi \rightarrow K$ as follows:

$$p(T) = \begin{cases} [(t, a)] & \text{if } \epsilon \leq t \leq \omega_a - \epsilon, \\ [(t, ba)] & \text{if } 0 \leq t \leq \epsilon, \\ [(t - \omega_a, ac)] & \text{if } \omega_a - \epsilon \leq t \leq \omega_a. \end{cases}$$

The map p is a continuous surjection, and there is a unique map $f : K \rightarrow K$ such that $f \circ p = p \circ \Phi$, where $\Phi : \mathcal{T}_\varphi \rightarrow \mathcal{T}_\varphi$ is the inflation and substitution homeomorphism. It is easy to verify that if $\varphi(a) = a_1 a_2 \dots a_k$, then $f(e_a) \subset e_{a_1} \cup e_{a_1 a_2} \cup e_{a_2} \cup \dots \cup e_{a_{k-1}} \cup e_{a_{k-1} a_k} \cup e_{a_k}$, and if $\varphi(a.b) = \dots c.d \dots$, then $f(e_{ab}) \subseteq e_c \cup e_{cd} \cup e_d$.

Lemma 2. $\varprojlim f \simeq \mathcal{T}_\varphi$.

Proof: The commuting diagram

$$\begin{array}{ccc} \mathcal{T}_\varphi & \xleftarrow{\Phi} & \mathcal{T}_\varphi \\ \downarrow p & & \downarrow p \\ K & \xleftarrow{f} & K \end{array}$$

induces a map $\hat{p} : \mathcal{T}_\varphi \rightarrow \varprojlim f$ given by $\hat{p}(T) = (p(T), p(\Phi^{-1}(T)), \dots)$.

Let $V := \{(\epsilon, a)\}_{a \in \mathcal{A}} \cup \{(\omega_a - \epsilon, a)\}_{a \in \mathcal{A}}$, and choose $\underline{x} = (x_0, x_1, \dots) \in \varprojlim f$. Suppose that for some $i \in \mathbb{N}$, $x_i \in V$. The choice of ϵ implies that $x_{i-1} \in e_b \setminus V$ for some $b \in \mathcal{A}$. That is, if $T \in \hat{p}^{-1}(\underline{x})$, then the 0^{th} tile of $\Phi^{1-i}(T)$, and the position of the origin within the interior of this tile, are determined, as are segments of T of approximate length $\lambda^i \omega_b$ about

the origin. It follows that $T \in \hat{p}^{-1}(\underline{x})$ is totally determined if $x_i \in V$ for arbitrarily large i .

If $x_i \notin V$ for arbitrarily large i , then for sufficiently large $n \in \mathbb{N}$, either $x_n \in e_{ab}$ for some $a, b \in \mathcal{A}$ or $x_n \in e_b \setminus V$ for some $b \in \mathcal{A}$. In the first case, the $(-1)^{st}$ and 0^{th} tiles of $\Phi^{1-n}(T)$, and the position of the origin within the interior of the union of these edges, are determined. In both cases, arguments identical to those above imply that arbitrarily large segments of T about the origin are determined, thus T is totally determined, and $\hat{p}$ is one-to-one. As p is onto, so is $\hat{p}$. $\square$

There are two essential ways in which the cohomology of the inverse limit of the natural map on the wedge of circles, that is, $\varinjlim A_\varphi^t$, may need to be modified to obtain the cohomology of the tiling space $\mathcal{T}_\varphi$.

First, if $\mathcal{T}_\varphi$ contains one or more cycles of asymptotic composants associated with a cycle of φ -periodic words of the form $\dots a_1.a_2\dots, \dots a_3.a_2\dots, \dots a_3.a_4\dots, \dots, \dots a_1.a_{2n}\dots$, those composants are ‘glued together’ in the inverse limit on the wedge of circles to form n -ods (possibly overlapping). The modified collaring described above, in particular, the construction of the subcomplex S , ‘pulls apart’ these cycles. The cohomology associated with the unglued cycles in $\mathcal{T}_\varphi$ is computed via $\check{H}^1(S)$ and must be added to $\varinjlim A^t$.

Second, there may be ‘extra’ cohomology appearing in $\varinjlim A^t$ that does not appear in $\mathcal{T}_\varphi$ that we identify below.

We find it easier to compute the cohomology of $\mathcal{T}_\varphi$ by working with a map g defined on K that is homotopic to f and leaves the transition edges invariant. We define g as follows.

First, let $\delta = \min\{\frac{\omega_a}{3}\}_{a \in \mathcal{A}}$. For $a \in \mathcal{A}$, define

$$h_a(x) = \begin{cases} \epsilon & \text{if } \epsilon \leq x \leq \delta, \\ \epsilon + \frac{\omega_a - 2\epsilon}{\omega_a - 2\delta}(x - \delta) & \text{if } \delta \leq x \leq \omega_a - \delta, \\ \omega_a - \epsilon & \text{if } \omega_a - \delta \leq x \leq \omega_a - \epsilon. \end{cases}$$

That is, h_a linearly maps the interval $[\delta, \omega_a - \delta]$ over $[\epsilon, \omega_a - \epsilon]$ and collapses the intervals $[\epsilon, \delta]$ and $[\omega_a - \delta, \omega_a - \epsilon]$ to $\{\epsilon\}$ and $\{\omega_a - \epsilon\}$, respectively.

Let $h : K \rightarrow K$ be defined as:

$$h(x, w) = \begin{cases} (h_a(x), w) & \text{if } x \in e_a, \text{ for } a \in \mathcal{A}, \\ (x, w) & \text{if } x \in e_{ab}, \text{ for } ab \in \mathcal{L}. \end{cases}$$

Finally, $g := h \circ f$. If $\varphi(a.b) = \dots c.d\dots$, then $g(e_{ab}) = e_{cd}$, hence $g(S) \subseteq S$.

We show below that the ‘extra’ piece of cohomology mentioned above may be computed as follows. We find the eventual range ER of $g|_S$, and the rank of the group generated by the coboundaries of the components of ER . Each

generator for this group is an eigenvector associated with eigenvalue 1 for a power of g^* acting on $H^1(K, S)$, and the rank of the group indicates the number of copies of $\mathbb{Z}$ to be quotiented out from $\varinjlim A^t$.

Whereas $\varprojlim f \simeq \mathcal{T}_\varphi$, $\varprojlim g$ is homeomorphic to a quotient of $\mathcal{T}_\varphi$ in which asymptotic composants for φ associated with φ -periodic words, should there be any, are glued in the direction of asymptoticity (each such pair of composants is associated with a pair of φ -periodic words of the form $\dots a.b\dots$, $\dots a.c\dots$, thus with the edges e_a , e_{ab} and e_{ac} , with the last two periodic under g). However, in this case, unlike the situation with the inverse limit space on the wedge of circles, gluing does not extend all the way to the origins of the associated tilings, and the next result says that this gluing does not change the cohomology of $\mathcal{T}_\varphi$.

Corollary 3. $\varinjlim g^* \simeq \check{H}(\mathcal{T}_\varphi)$.

Proof: Since f and g are homotopic, $\varinjlim g^* = \varinjlim f^* \simeq \check{H}(\mathcal{T}_\varphi)$. $\square$

Denote the eventual range $\cap_{n \geq 0} g^n(S)$ of $g|_S$ by ER . Let k be the number of connected components of ER and l the number of independent 1-cycles in ER ; we call l the *asymptotic cycle rank* of φ . (Note that if φ is proper, ER consists of a single edge e_{ab} . If φ forces its border, then each component of ER is a single edge.)

Our main result is the following:

Theorem 4. $\check{H}^1(\mathcal{T}_\varphi) \simeq \mathbb{Z}^l \oplus \varinjlim A^t / G$, where $G \simeq \mathbb{Z}^{k-1}$.

Proof: There is an $m \in \mathbb{N}$ such that $g^m(C) = C$ for each component C of ER and $g^m(S) \subseteq ER$. Pick a point c from each component C of S , and let $\hat{c}$ denote the corresponding dual generator in $H^0(S)$. For each component C of ER , the element

$$v_C = \Sigma \{ \hat{c}' : C' \text{ is a component of } S, g^m(C') \subseteq C \}$$

is fixed by $(g^*)^m : H^0(S) \rightarrow H^0(S)$. Furthermore, the eventual range of g^* has basis $\{v_C : C \text{ a component of } ER\}$, and the sum of these basis elements, Σv_C , generates the range η of the augmentation map. Thus, choosing $k-1$ of the components, $C_1, \dots, C_{k-1}$, of ER , the cosets $\{v_{C_1} + \eta, \dots, v_{C_{k-1}} + \eta\}$ form a basis for the eventual range of g^* on the reduced cohomology $\tilde{H}^0(S) := H^0(S)/\eta$, and each of these basis elements is an eigenvector for $(g^*)^m$ with eigenvalue 1.

It is clear that g^m is homotopic to the identity on ER , so that $\varinjlim g^* : H^1(S) \rightarrow H^1(S) \simeq \mathbb{Z}^l$, where l is the asymptotic cycle rank of φ . To see how the 1-cycles of ER contribute to the cohomology of $\mathcal{T}_\varphi$, consider the morphism of the exact sequence of the pair (K, S) induced by $g : (K, S) \rightarrow (K, S)$:

$$\begin{array}{ccccccc}
0 & \rightarrow & \tilde{H}^0(S) & \xrightarrow{\delta} & H^1(K, S) & \rightarrow & H^1(K) \rightarrow H^1(S) \rightarrow 0 \\
& & \downarrow g_0^* & & \downarrow g_1^* & & \downarrow g_2^* & \downarrow g_3^* \\
0 & \rightarrow & \tilde{H}^0(S) & \xrightarrow{\delta} & H^1(K, S) & \rightarrow & H^1(K) \rightarrow H^1(S) \rightarrow 0
\end{array}$$

Taking direct limits, we get an exact sequence

$$0 \rightarrow G_0 \xrightarrow{\vec{\delta}} G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$$

in which $G_1 \simeq \varinjlim A^t$, $G_2 := \varinjlim g_2^* \simeq \check{H}^1(\mathcal{T}_\varphi)$, $G_3 := \varinjlim g_3^* \simeq \mathbb{Z}^l$, and $G_0 := \varinjlim g_0^* \simeq \mathbb{Z}^{k-1}$ as described above. The short exact sequence

$$0 \rightarrow \operatorname{coker}(\vec{\delta}) \rightarrow G_2 \rightarrow G_3 \rightarrow 0$$

splits to give $\check{H}^1(\mathcal{T}_\varphi) \simeq \operatorname{coker}(\vec{\delta}) \oplus \mathbb{Z}^l$.

Since $H^1(K)$ is free, so is $\operatorname{coker}(\delta)$. In particular, the image under δ of a basis of $\tilde{H}^0(S)$,

$\{\delta(\hat{c}_1 + \eta), \dots, \delta(\hat{c}_{k-1} + \eta)\} \cup \{\delta(\hat{c} + \eta) : C \text{ a component of } S, C \cap ER = \emptyset\}$,

extends to a basis for $H^1(K, S)$ with respect to which $(g_1^*)^m$ takes the form

$\begin{pmatrix} I & 0 & E_2 \\ E_1 & 0 & E_3 \\ 0 & 0 & A_1 \end{pmatrix}$, where I is the $(k-1) \times (k-1)$ identity matrix, and the

diagonal 0 matrix is size $j \times j$, where $j := (\# \text{ of components in } S) - k$. Note that in the basis $\{\hat{e}_a : a \in \mathcal{A}\}$, g_1^* is represented by A^t , the transpose of the

transition matrix for φ . Thus $\begin{pmatrix} I & 0 & E_2 \\ E_1 & 0 & E_3 \\ 0 & 0 & A_1 \end{pmatrix}$ is unimodularly equivalent

to $(A_1^t)^m$, and $\operatorname{coker}(\vec{\delta}) \simeq \varinjlim (\overline{g_1^*})^m$, where $\overline{g_1^*} : \operatorname{coker}(\delta) \rightarrow \operatorname{coker}(\delta)$ is given by $\overline{g_1^*}(x + \delta(\tilde{H}^0(S))) = g_1^*(x) + \delta(\tilde{H}^0(S))$. Thus $\operatorname{coker}(\vec{\delta}) \simeq \varinjlim A_1$ and we have $\check{H}^1(\mathcal{T}_\varphi) \simeq \varinjlim A_1 \oplus \mathbb{Z}^l$. $\square$

Note: $\varinjlim \begin{pmatrix} I & 0 \\ E_1 & 0 \end{pmatrix} \simeq \mathbb{Z}^{k-1}$. There are conditions that imply that $\varinjlim A^t$ splits as $\mathbb{Z}^{k-1} \oplus \varinjlim A_1$. For instance, if there is a matrix B so that $BA_1 + \begin{pmatrix} I & 0 \\ E_1 & 0 \end{pmatrix} = -\begin{pmatrix} E_1 \\ E_2 \end{pmatrix}$, then $\varinjlim A^t$ splits as described. We do not know if there is always such a splitting.

We rephrase the above discussion in a way that is more consistent with actual computations (for instance, see Example 8) and allows for a more precise statement of Theorem 4.

Let $f_i = (0 \dots 010 \dots)^t$ be the i^{th} standard basis vector for $\mathbb{R}^d$ and $C_1, \dots, C_p$ the components of the complex S . For $i = 1, \dots, p-1$, let

$w_i := \sum f_j - \sum f_m$, where the first sum is over those j for which $e_j = [\epsilon, \lambda_j - \epsilon] \times \{j\}$ terminates in C_i (i.e., $(\lambda_j - \epsilon, j) \in C_i$) and the second sum is over those m for which e_m originates in C_i (i.e., $(\epsilon, m) \in C_i$). Then $\{w_1, \dots, w_{p-1}\}$ extends to a basis $\{w_1, \dots, w_{p-1}, w_p, \dots, w_d\}$ of $\mathbb{Z}^d$. Let P be the matrix with columns $\{w_1, \dots, w_{p-1}, w_p, \dots, w_d\}$. Then $P^{-1}A^tP$ has the form $\begin{pmatrix} E & F \\ 0 & A_1 \end{pmatrix}$ in which A_1 is size $(d - p + 1) \times (d - p + 1)$.

Theorem 4'. Let φ be an aperiodic primitive substitution with asymptotic cycle rank l and A_1 as above. Then $\check{H}^1(\mathcal{T}_\varphi) \simeq \varinjlim A_1 \oplus \mathbb{Z}^l$.


Corollary 5. Let φ , K , and S be as above. If $\check{H}^*(S) \simeq 0$, then $\check{H}^1(\mathcal{T}_\varphi) \simeq \varinjlim A_\varphi^t$.

Example 6. *The Fibonacci substitution:*

If φ is the Fibonacci substitution ($\varphi(1) = 12, \varphi(2) = 1$) (see Example 1), then S has a single connected component, and the eventual range of $g|_S$ is the connected pair of edges $ER = e_{11} \cup e_{21}$. In this case, $\check{H}^0(S) \simeq 0 \simeq H^1(S)$, and $\check{H}^1(\mathcal{T}_\varphi) \simeq \varinjlim \left[\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^t \right] \simeq \mathbb{Z}^2$.

Example 7. *The Morse-Thue substitution:*

Let φ denote the Morse-Thue substitution ($\varphi(1) = 12, \varphi(2) = 21$).



In this case, all transitions are allowed, and K is as above. S has a single connected component, $ER = S$, $\check{H}^0(S) \simeq 0$ and $H^1(S) = \mathbb{Z}$. Then $\check{H}^1(\mathcal{T}_\varphi) \simeq \varinjlim \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \oplus \mathbb{Z} \simeq \mathbb{Z}[1/2] \oplus \mathbb{Z}$.

Example 8. *Disconnected S :*

Let φ be the substitution: $\varphi(1) = 12341$, $\varphi(2) = 12$, $\varphi(3) = 3423$, $\varphi(4) = 42$. The set S has two connected components associated with the two sets of allowed transitions: $\{11, 41, 21, 23, 12, 42\}$ and $\{34\}$. The set ER consists of two connected components associated with $\{21, 11, 23\}$ and $\{34\}$. Then $\check{H}^0(S) \simeq \mathbb{Z}$ has basis $\{\hat{c}_1\}$, where $\hat{c}_1$ is the dual generator of a vertex c_1 in e_{34} . It follows that $\{\delta\hat{c}_1 = \hat{e}_3 - \hat{e}_4\}$ extends to the basis $\{\hat{e}_3 - \hat{e}_4, \hat{e}_1, \hat{e}_2, \hat{e}_3\}$

of $H^1(K, S)$ with respect to which g_1^* has matrix $\begin{pmatrix} 1 & 0 & -1 & 0 \\ 0 & 2 & 1 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 2 & 2 \end{pmatrix}$. Since

there are no 1-cycles in S , $\check{H}^1(\mathcal{T}_\varphi) \simeq \varinjlim \begin{pmatrix} 2 & 1 & 1 \\ 1 & 1 & 0 \\ 0 & 2 & 2 \end{pmatrix}$.

REFERENCES

- [AP] J.E. Anderson and I.F. Putnam, Topological invariants for substitution tilings and their associated C^* -algebras, *Ergodic Theory & Dynamical Systems* **18** (1998), 509-537.
- [BD] M. Barge and B. Diamond, A complete invariant for the topology of one-dimensional substitution tiling spaces, *Ergodic Theory and Dynamical Systems* **21** (2001), 1333-1358.
- [BJV] M. Barge, J. Jacklitch, and G. Vago, Homeomorphisms of one-dimensional inverse limits with applications to substitution tilings, unstable manifolds, and tent maps, *Contemporary Mathematics* **246**, 1 - 15.
- [BJKR] O. Bratteli, P. Jorgenson, K. Kim, and F. Roush, Decidability of the isomorphism problem for stationary AF-algebras, *Ergodic Theory & Dynamical Systems* **21**(2001), 1625-1655.
- [Dur] F. Durand, A characterization of substitutive sequences using return words, *Discrete Math* **179**(1998), 89-101.
- [Mo] B. Mossé, Puissances de mots et reconnaissabilité des points fixes d'une substitution, *Theoretical Computer Science* **99** (1992), 327-334.
- [So] B. Solomyak, Nonperiodicity implies unique composition for self-similar translationally finite tilings, *Discrete Comput. Geometry* **20** (1998), 265-279.

Department of Mathematics, Montana State University, Bozeman, MT 59717
 barge@math.montana.edu
 Department of Mathematics, College of Charleston, Charleston, SC 29424
 diamondb@cofc.edu